

Dietary protein content for an optimal diet: a clinical view

Lidia Santarpia*, Franco Contaldo & Fabrizio Pasanisi

Interuniversity Centre for Obesity and Eating Disorders, Internal Medicine and Clinical Nutrition, Department of Clinical Medicine and Surgery, Federico II University, Naples, Italy

Abstract

The dietary protein role in different clinical nutritional conditions and some physio-pathological perspectives is a current and hot topic to discuss. Recent Proceedings of the Protein Summit 2, joining more than 60 nutrition scientists, health experts, and nutrition educators, suggest to increase plant but, in particular, animal protein intake because richer in leucine and consequently more effective to influence anabolic protein metabolism. The Panel conclusions are in apparent contradiction with the nutritional ecology statements, which strongly sustain the reduction of animal origin foods in the human diet and are currently concerned about the excessive, mainly animal protein intake in western and westernized Countries. In conclusion, it is time to carefully evaluate protein and aminoacid intake accurately considering quality, digestibility, daily distribution and individual characteristics.

Keywords dietary protein; animal proteins; essential aminoacids; leucine

Received: 30 September 2016; Accepted: 13 November 2016

**Correspondence to: Lidia Santarpia, Department of Clinical Medicine and Surgery, Federico II University, Via Pansini, 5 80131 Naples, Italy. Tel: 0039 081 746 2333; Fax: 0039 081 746 2333, Email: lidia.santarpia@unina.it*

The American Journal of Clinical Nutrition (AJCN) recently published, as supplement,^{1–6} the Proceedings of the Protein Summit 2 to evaluate the dietary protein role in various clinical nutritional conditions and some physio-pathological perspectives such as weight management, metabolic activity, healthy aging, and healthier diets within energy balance. The Summit, as well as AJCN supplement, joining more than 60 nutrition scientists, health experts, and nutrition educators, was supported also by several ‘meat companies’.¹

To summarize the conclusions, and starting from weight management (optimal weight loss) in obese patients, the Panel¹ showed a greater weight loss, fat mass loss, better preservation of lean body mass, a more significant reduction of blood pressure, serum tryglicerides, and waist circumference following higher protein vs. low protein restricted diets, both in the short and long term. A mild effect on satiety, but not on satiation, was also disclosed. Overall data suggest that protein content of restrictive diets should be in the range of 1.2–1.6 g protein/kg/body weight (BW)/day with the general indication to take 20–30 g proteins at each main meal

(breakfast, lunch, and dinner). These findings have been confirmed in a recent paper by Wejis and Wolfe⁷ in a sample of obese older adults (over 55 yrs), showing that protein requirement during weight loss should be of at least 1.2 g/kg BW and 1.9 g/kg fat-free mass to obtain a satisfactory muscle mass accretion.

As far as the proteins’ metabolic role, the panel states that recommended dietary allowances are based on nitrogen balance (NB) studies, which simply reflect overall nitrogen retention under conditions of energy balance in healthy young adults.⁸ Furthermore, dietary protein intake is calculated on the minimum amount to reach NB and the minimum aminoacid (AA) requirements. On the other hand, a more accurate and realistic approach to protein balance is to evaluate the essential or indispensable AA requirements to obtain metabolic benefits, and not only maintenance, as improved glycemic regulation, recovery after trauma, improved lean body mass function, etc. The Panel suggests the Indicator Amino Acid Oxidation (IAAO) method, as an alternate approach to NB, IAAO method, emphasizes that if

indispensable AAs are deficient in the diet, all other AAs will be oxidized, according to the different physio-pathological conditions.^{9,10} As a matter of the fact, the IAAO method suggests, once again, a protein requirement of 1.2 g/kg/BW/day, well higher than the current recommendations of dietary guidelines.⁸

Other studies^{3,11,12} identify AAs themselves as potential signals to activate protein synthesis and not only as substrates for protein metabolism. Among various signalling systems, the leucine-induced activation mammalian Target of Rapamycin Complex 1 (mTORC-1) pathway has a main role, in particular on muscle protein synthesis: the leucine-induced activation of mTORC1 leads to the stimulation of skeletal muscle synthesis, preferentially within 2 h after the ingestion of a meal containing at least 20–30 g leucine rich proteins.^{13–15} This response, called ‘meal threshold’, reinforced by regular physical exercise, decreases by aging, thus suggesting to slightly increase protein intake in the elderly, a quite new nutritional education suggestion, besides the traditional indication to maintain a regular physical exercise to prevent sarcopenia¹⁶ through a positive effect on mTORC1 pathway too.¹⁷ Although an increase of both plant and animal protein intake is suggested,¹⁸ the Panel remarks that animal proteins are richer in leucine and consequently more effective to influence anabolic protein metabolism. A warning is also risen on the current dietary recommendation to strongly reduce saturated and solid fat intake, which in dietary practice means to reduce animal proteins and related food groups. The chronic reduction in animal-based¹⁹ food groups might have, as secondary effect, the reduced intake of some ‘nutrients of concern’ (i.e. nutrients with a daily intake generally lower than recommended) as Calcium, vitamin D, Potassium, Iron, Folate for animal-based protein foods^{18,19} and Dietary Fiber, vitamin E, and Magnesium for plant-based protein foods.^{18,20}

The Panel finally affords the topic of translating current scientific evidences in the clinical practice; in other words, do we need to reevaluate the current dietary guidelines?

The Panel¹ speculates that the current recommended dietary allowances of 0.8 g Proteins/kg BW/day are based on Estimated Average Requirement of 0.66 g/kg BW/day, the ‘average daily nutrient intake level estimated to meet the requirement of half the healthy individuals in a particular life stage and gender group’.^{20,21} In conclusion, 0.8 g proteins/kg BW/day seem to be too low, at least in some clinical conditions and the NB method inaccurate to properly monitor individual protein requirements. Consequently, the Panel proposes ‘flexibility’ also for protein intake in order to create a variety of eating planes for single individuals.

In nutritional terms, the Panel supports the recommendation of Acceptable Macronutrient Distribution Ranges (AMDR), which aims to maintain adequate macronutrient (and micronutrient) intakes against the backdrop of adequate energy intake. According to the Institute of Medicine,²¹ the AMDR for Carbohydrates ranges between 45% and 65%, Fat

20% and 35%, and Protein 10% and 35% of total energy intake.

A convincing statement by the Panel, at least in our opinion, is the timing of protein intake to be equally distributed at a minimum quantity of 20–30 g/meal during the three main courses.

Emphasis is also given to proteins’ AA composition; strong recommendation is finally given to increase the recommended protein intake from 0.8 to 1.0–1.2 g/kg/day, at least in the elderly and in some other physio-pathological conditions.

These conclusions reached after reading the AJCN supplement are in apparent contradiction with the nutritional ecology statements, which (actually since more than 10 years) strongly sustain the reduction of animal origin foods in the human diet and are currently concerned about the excessive (mainly animal) protein intake in western and westernized Countries.^{22,23} Recent observations support the suspect that high dietary protein intake (when protein replace carbohydrates) is associated, in the general population, to a greater risk of weight gain.²⁴

The amount (and quality) of protein intake is still a ‘hot’ nutritional topic, to widely debate and clarify.

It is out of the interest of this paper to discuss on dietary protein quality; nevertheless, Food and Agriculture Organization (FAO) has recently reconsidered the use of a new score to quantify dietary protein quality: the Digestible, Indispensable (or Essential) Aminoacid score, and DIAAS.²⁵

This score highlights IAA content and protein digestibility as indicator of protein biological value, a factor of valuable clinical and physiological interest.^{26,27}

As a preliminary observation, it appears necessary a large consensus on what does g protein/kg BW/day mean. What do we intend for BW: actual, ideal BW, or what else? How to treat severely underweight or overweight, young or elderly, sarcopenic or body builder individuals? The suggestion of 0.8 or 1.2 or other amounts of dietary protein/kg BW weight has a marked bias because of the variability in body composition, clearly not identifiable by the simple measurement of BW and body mass index! We would suggest, and actually we do, at least in the current clinical practice but not in the critically ill patient, to adopt as reference BW, the ‘desirable BW’, calculated from the mean value of body mass index between 18.5 and 25 kg/m². Nevertheless, we ask a discussion on this question. Furthermore, adequate protein intake requires a specific evaluation for critically ill patients, in particular when undergoing artificial nutrition. We also consider that, still confirming for the general population the current dietary recommendation of 0.8 g protein/kg ‘desirable’ BW/day, AMDR could be the reasonable dietetic approach in the clinical practice but also a rationale for dietary guidelines. It allows some flexibility in the dietary prescription, given the extreme heterogeneity of populations with large ethnic, cultural, social, economic inter-individual differences and peculiarities, wider age range with a marked

increase of elderly, over 65 yrs, people, a significant decrease in physical activity at all ages, etc. General recommendations to pursue a more vegetarian, Mediterranean style, ecologically compatible diet should remain, provided that a variety in consumption of high nutritional value, plant and animal, foods is assured. A more accurate protein-rich food distribution at the main three meals (in particular by increasing protein intake at breakfast, at least in the Mediterranean Countries with a simultaneous reduction at dinner time) could improve protein balance and consequently reduce protein intake. A protein sparing effect will be also obtained by a regular physical exercise. A higher intake of mixed plant and animal proteins than the current 0.8g/kg BW/day may be suggested, associated to regular physical exercise, in restrictive diets for obese patients and in (otherwise healthy) sarcopenic elderly individuals.

In our opinion, the topic of protein intake recommendations has not been exhaustively examined so far at least in the clinical nutritional practice: it is now time to carefully evaluate protein and AAs intake as usually carried out for

fat and CHO intakes accurately considering quantity but also quality, digestibility, timing of daily distribution, composition of the meal, individual fat-free mass characteristics, total daily energy intake, concomitant diseases, etc.

A special attention is also to pay to 'nutrients of concern' whose assumption may be easily affected by the composition of the prescribed diets, including protein rich foods.

Acknowledgement

The authors certify that they comply with the ethical guidelines for publishing in the Journal of Cachexia, Sarcopenia, and Muscle: update 2015.²⁸

Conflict of interest statement

None declared.

References

- Rodriguez NR. Introduction to Protein Summit 2.0: meal requirements for protein to optimize metabolic roles of aminoacids, continued exploration of the impact of high-quality protein on optimal health. *Am J Clin Nutr* 2015;**101**:1317S–1319S.
- Leidy HJ, Clifton PM, Astrup A, Wycherley TP, Westerterp-Plantenga MS, Luscombe-Marsh ND, Woods SC, Mattes RD. The role of protein in weight loss and maintenance. *Am J Clin Nutr* 2015;**101**:1320S–1329S.
- Layman DK, Anthony TG, Rasmussen BB, Adams SH, Lynch CJ, Brinkworth GD, Davis TA. Defining meal requirements for protein to optimize metabolic roles of aminoacids. *Am J Clin Nutr* 2015;**101**:1330S–1338S.
- Paddon-Jones D, Campbell WW, Jacques PF, Kritchevsky SB, Moore LL, Rodriguez RR, van Loon JC. Protein and healthy aging. *Am J Clin Nutr* 2015;**101**:1339S–1345S.
- Phillips SM, Fulgoni VL III, Heaney RP, Nicklas TA, Slavin JL, Weaver CM. Commonly consumed protein foods contribute to nutrient intake, diet quality, and nutrient adequacy. *Am J Clin Nutr* 2015;**101**:1346S–1352S.
- Rodriguez NR, Miller SL. Effective translation of current dietary guidance: understanding and communicating the concepts of minimal and optimal levels of dietary protein. *Am J Clin Nutr* 2015;**101**:1353S–1358S.
- Wejis PJM, Wolfe RR. Exploration of the protein requirement during weight loss in obese older adults. *Clin Nutr* 2016;**35**:394–398.
- Institute of Medicine, Food and Nutrition Board. *Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fatty Acids, Cholesterol, Protein and Amino Acids*. Washington (DC): National Academies Press; 2002.
- Munro H, Crim M. The protein and aminoacids. In Shils ME, Young VR, eds. *Modern Nutrition in Health and Disease*, 7th ed. Philadelphia: Lea and Febiger; 1988. p1–37.
- Millward DJ. An adaptive metabolic demand model for protein and aminoacid requirements. *Br J Nutr* 2003;**90**:249–260.
- Lynch CJ, Halle B, Fujii H, Vary T, Wallin R, Damuni Z, Hutson S. Potential role of leucine metabolism in the leucine-signaling pathway involving mTOR. *Am J Physiol Endocrinol Metab* 2003;**285**:E854–E863.
- Li F, Yin Y, Kong X, Wu G. Leucine nutrition in animals and humans: mTOR signaling and beyond. *Amino Acids* 2011;**41**:1185–1193.
- Paddon-Jones D, Rasmussen BB. Dietary protein recommendations and the prevention of sarcopenia. *Curr Opin Clin Nutr Metab Care* 2009;**12**:86–90.
- Layman DK. Dietary guidelines should reflect new understandings about adult protein needs. *Nutr Metab* 2009;**6**: doi: 10.1186/1743-7075-6-12.
- Hartman JW, Tang J, Wilkinson S, Tarnopolsky M, Lawrence R, Fullerton A, Phillips S. Consumption of fat-free fluid milk after resistance exercise promotes greater lean mass accretion than does consumption of soy or carbohydrate in young, novice, male weight-lifters. *Am J Clin Nutr* 2007;**86**:373–381.
- Breen L, Phillips SM. Interactions between exercise and nutrition to prevent muscle waste during aging. *Br J Clin Pharmacol* 2013;**75**:708–715.
- Yang Y, Breen L, Burd N, Hector AJ, Churchward-Venne TA, Josse AR, Tarnopolsky MA, Phillips SM. Resistance exercise enhances myofibrillar protein synthesis with graded intakes of whey proteins in older men. *Br J Nutr* 2012;**108**:1780–1788.
- Huth PJ, Fulgoni VL, Keast DR, Park KI, Auestad N. Major food sources of calories, added sugars, and saturated fat and their contribution to essential nutrients intake in the U.S. diet: data from the national health and nutrition examination survey (2003-2006). *Nutr J* 2013;**12**:–116.
- Murphy MM, Spungen JH, Bi X, Barraj LM. Fresh and fresh lean pork are substantial sources of key nutrients when these products are consumed by adults in the United States. *Nutr Res* 2011;**31**:776–783.
- O'Neil CE, Keast DR, Fulgoni VL, Nicklas TA. Tree nut consumption improves nutrient intake and diet quality in US adults: an analysis of National Health and Nutrition Examination Survey 1999–2004. *Nutr Res* 2012;**32**:185–194.
- Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fatty Acids, Cholesterol, Protein and Amino Acids*. Washington (DC): National Academies Press; 2002.

22. Institute of medicine, dietary reference intakes for energy, carbohydrate, fiber, fat, protein, and amino acids (macronutrients). Washington (DC): The National Academies Press; 2002-2005.
23. Pimentel D, Pimentel M. Sustainability of meat-based and plant-based diets and the environment. *Am J Clin Nutr* 2003;**18**:660S–663S.
24. Contaldo F, Pasanisi F. High protein diet, obesity and the environment. *Am J Clin Nutr* 2006;**83**:387.
25. Food and Agriculture Organization of the United Nations/WHO. Dietary protein quality evaluation in human nutrition: report of an FAO expert consultation, Rome, Italy 2013, FAO Food and Nutrition Paper 92
26. Wolfe RR, Rutherford SM, Kim IY, Moughan PJ. Protein quality as determined by the Digestible Indispensable Amino Acid Score: evaluation of factors underlying the calculation. *Nutr Rev* 2016;**74**:584–599.
27. Alonso PH, Salas-Salvado J, Canela MR and other 13 authors. High dietary protein intake is associated with an increased body weight and total death risk. *Clin Nutr* 2016;**35**:496–506.
28. von Haehling S, Morley JE, Coats AJS, Anker SD. Ethical guidelines for publishing in the Journal of Cachexia, Sarcopenia and Muscle: update 2015. *J Cachexia Sarcopenia Muscle* 2015;**6**:315–316.